

MODULAR VECTOR SYSTEMS

Priority Claim and Related Applications

0
1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213
2214
2215
2216
2217
2218
2219
2220
2221
2222
2223
2224
2225
2226
2227
2228
2229
22

0
1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20

To give but a few examples, plasmid vectors that replicate in different hosts, with different copy numbers, have been prepared (e.g., bacterial vectors designed to have either relaxed or stringent control of replication; yeast vectors with either a 2 μ or centromeric replication origin, mammalian vectors containing viral [e.g., SV40 or BPV] origins of replication, etc.). Vectors have been engineered to allow ready detection of insertion events (e.g., by creation or disruption of a selectable or detectable marker), to direct high levels of expression of proteins encoded by inserted sequences (e.g., under the control of transcription, splicing, and/or translation signals active in a given host system), to generate gene fusions that allow analysis of expression of inserted sequences (e.g., by analysis of B-galactosidase, chloramphenicol transferase, luciferase, or green fluorescent protein activity, etc.), or to create fusion proteins with experimentally useful attributes (e.g., easy purification, desired cellular localization, etc.). Vectors have been designed that are particularly useful for determining the sequence of inserted fragments (e.g., by allowing easy production of single-stranded DNA), or for producing RNA (sense or antisense) from the inserted sequences. Most companies that sell molecular biology reagents include among their products vectors that they have developed to be particularly useful for designated applications (see, for example, catalogs provided by Amersham Pharmacia Biotech, Piscataway, NJ; Promega Corporation, Madison, WI; Invitrogen Inc., Carlsbad, CA; Life Technologies, Inc., Rockville, MD; New England Biolabs, Beverly, MA; Stratagene, Inc., La Jolla, CA).

Of course, the universe of genetic "vectors" is not limited to circular molecules derived from bacterial plasmids. Any nucleic acid molecule that includes sequences sufficient to direct *in vivo* or *in vitro* self-replication can be employed as a vector.

Typically, such replication sequences include a replication origin that directs duplication of the vector sequence in a host system (typically a transformed cell). Alternatively, sequences that direct integration of the vector into another nucleic acid molecule that is present in and replicated by the relevant host system can be sufficient to achieve vector
5 (and insert) replication.

Most vectors in use today are derived from naturally-occurring bacterial plasmids, bacteriophages, or other viruses. Some vectors contain features of more than one of these systems. Almost all of the commonly-used vectors contain one or more restriction sites designed for convenient insertion of fragments; most have at least one polylinker (see, for
10 example, the vector database maintained at <http://vectorbd.atcg.com/vectordb/vector.html>, the contents of which as of July 19, 2000 are included herein as Appendix A).

Despite the broad availability of vectors from commercial and other sources, each one has features selected by the relevant manufacturer rather than the experimental user.
15 It is not uncommon for a researcher to have to modify an available vector to suit his experimental needs, or alternatively to modify his experimental design to accommodate the available vectors. There remains a need for the development of techniques and reagents that would allow a researcher to readily design and assemble vector(s) appropriate to his experimental needs.

Summary of the Invention

The present invention encompasses the recognition that vectors are comprised of modular elements and need not be provided as discrete nucleic acid molecules into which

fragments of interest are inserted. Rather, vectors can themselves be assembled from pieces that contain part or all of individual useful elements. In certain preferred embodiments of the invention, fragments corresponding to pieces of what is traditionally viewed as the "vector backbone" are provided individually and are linked to one another

5 substantially simultaneously with the linkage that associates vector sequences with insert sequences.

According to the present invention, components of a vector can be defined as one of a variety of categories of vector elements. For example, sequences that allow the vector to replicate in a host system may be classified as "replication elements".

10 Similarly, sequences that allow host cells containing a vector to survive experimental conditions that kill otherwise identical host cells lacking a vector may be classified as "replication elements"; sequences that allow detection but not selection of host cells containing vector sequences, or host cells containing vector and insert sequences, may be classified as "detectable elements"; sequences that can act to direct expression (i.e.,

15 transcription, splicing, and/or translation) of other sequences can be classified as "expression elements". Other categories of elements may also be defined as discussed in further detail herein.

The present invention allows a researcher to select individual elements from one or more categories of vector elements, and to combine the selected element(s) with one or

20 more individual element(s) with one another to assemble vectors that contain a desired collection and arrangement of elements. Individual vector elements, or portions or combinations thereof, are provided on separate "vector fragments" that are linked together to create the final vector. Thus, the present invention provides techniques and

reagents useful in the assembly of vectors from individual vector fragments. Preferably, a vector assembled according to the present invention will include at least a replication element. More preferably, the vector will include one or more additional elements selected from the group consisting of additional replication elements (e.g., effective in
5 different host systems), selectable markers, detectable markers, expression elements, fusion protein elements, mobile elements, recombination elements, cleavage site elements, etc. The inventive techniques and reagents may be employed to link two or more vector fragments to one another, serially or simultaneously, and also to link vector fragments with one or more insert fragments (again, serially or simultaneously).

10 In particularly preferred embodiments of the present invention, one or more of the vector and insert fragments used in the assembly of a final hybrid construct is prepared without the use of restriction enzymes (or any endonuclease). Most preferably, substantially all of the fragments that become linked together to produce a final assembled molecule are prepared without the use of restriction enzymes. In particularly
15 preferred embodiments of the invention, RNA-Overhang Cloning and/or DNA Overhang Cloning are employed to produce vector and/or insert fragments. Also, in certain preferred embodiments of the invention, vector fragments, and optionally insert fragments, are linked to one another by ligation-independent cloning (i.e., without the use of a ligase enzyme).

Description of the Drawing

Figure 1 depicts assembly of a hybrid molecule comprising λ vector elements and an insert, according to the present invention.

Figure 2 shows assembly of a hybrid molecule comprising bacterial vector elements and an insert in a three-molecule linkage reaction according to the present invention.

Figure 3 depicts assembly of a hybrid molecule containing bacterial vector
5 elements and an insert according to the present invention. Two vector fragments and one insert fragment are linked together to form a hybrid that can be selected by growth in the presence of tetracycline and lack of growth in the presence of ampicillin.

Figure 4 depicts assembly of a hybrid molecule comprising bacterial vector
10 elements and an insert according to the present invention. Two vector fragments, each of which contains a portion of a detectable element, and one insert fragment are linked together to form a hybrid. Hybrids that contain insert can be distinguished from those that do not by a blue/white screen.

Figure 5 shows assembly of a hybrid molecule containing bacterial vector
15 elements and an insert according to the present invention. Two vector fragments, one of which contains a bacterial origin of replication and a first portion of a LacZ gene and one of which contains an ampicillin resistance gene and a second portion of the LacZ gene are linked to an insert fragment. Hybrids can be selected by growth in the presence of ampicillin; those containing insert can be distinguished from those lacking insert by a blue/white screen.

20 Figure 6 shows assembly of a hybrid molecule from three vector fragments and one insert fragment. Linkage of the four fragments re-creates two vector elements, and operatively links a third (the promoter) with the insert sequences.

Figure 7 shows collections of vector fragments, each of which contains only a single vector element, that may alternatively be linked to each other and an insert to form a hybrid molecule according to the present invention.

Figure 8 depicts a kit comprising two collections of vector fragments that can be used in various combinations to create vectors with different attributes according to the present invention. The first collection of vector fragments contains three fragments, each of which includes the pGal promoter and a first portion of a selectable marker selected from the group consisting of the *URA3*, *TRP1*, and *HIS3* genes. The second collection of vector fragments contains six different fragments, each of which contains a second portion of one of the selectable markers, and an origin of replication that is either a centromeric origin or a 2 μ origin.

Figure 9 depicts assembly of a hybrid molecule from two vector fragments and one insert fragment, each of which was prepared by DOC, according to the present invention. Panel A shows the generation of the two vector fragments; Panel B depicts the ligation of these two fragments with the insert fragment to produce the final hybrid.

Figure 10 shows a hybrid molecule assembled from two vector fragments and an insert fragment, each of which was prepared by DOC, according to the present invention.

Figure 11 shows the primers used (3NT5'OST [SEQ ID NO:___]; 3NT3'OHT [SEQ ID NO:___]; 3NT5'KHT [SEQ ID NO:___]; 3NT3'KST [SEQ ID NO:___]; INT5'OSI [SEQ ID NO:___]; INT3' Ori(s) [SEQ ID NO:6]; INT5'KAN [SEQ ID NO:11]; INT3'KAN [SEQ ID NO:12].

Definitions

"Element"-- The term "element" is used herein to refer to a region of nucleic acid sequence that imparts a particular functional or structural characteristic upon the molecule.

"Expression"-- "Expression" of a nucleic acid sequence, as that term is used
5 herein, refers to one or more of the following events: (a) production of an RNA template from a DNA sequence (e.g., by transcription); (b) processing of an RNA transcript (e.g., by splicing, editing, and/or 3' end formation); (c) translation of an RNA has been into a polypeptide or protein; (d) post-translational modification of a polypeptide or protein.

10 "Fragment"-- A "fragment", as that term is used herein, is an individual nucleic acid molecule that can be hybridized or linked with one or more other fragment molecules to produce a hybrid molecule. Preferably, a fragment contains at least a portion of a selected sequence element so that, when the fragments are linked together, a hybrid molecule is generated that contains a predetermined collection and arrangement of sequence elements. In certain preferred embodiments of the invention, each fragment
15 contains at least one intact sequence element. In other preferred embodiments, each fragment contains only one intact sequence element. In still other preferred embodiments, at least one fragment contains only a portion of a particular sequence element (though the fragment may also contain a complete copy of a different sequence element). Preferably, that fragment will become linked with another fragment so that the
20 complete sequence element is reassembled in the final hybrid. Alternatively or additionally, fragments are selected so that different hybrid molecules can be produced from linkage of the same collection of fragments, and such different hybrids can be distinguished from one another on the basis of whether a particular sequence element is

recreated in the hybrid. Preferred fragments for use in accordance with the present invention are prepared without the use of restriction enzymes. Most preferably they are prepared by polymerase chain reaction (PCR) amplification according to ROC or DOC techniques (see, for example, USSN 60/114,909, USSN 09/225,990, and Coljee et al.,
5 *Nature Biotechnology* 18:789, July 2000, each of which is incorporated herein by reference in its entirety). Preferred fragments are double stranded nucleic acid molecules with at least one single-stranded overhang.

"Host system"-- A "host system" according to the present invention is any *in vivo* or *in vitro* system into which a vector is introduced. Preferably, the host system is a cell
10 or organism. Any type of cell, including a bacterial cell, yeast cell, plant cell, or animal cell, can be a host cell. Cells in culture and cells that are part of living tissues or organisms can also be host cells.

"Hybrid"-- A "hybrid" nucleic acid molecule according to the present invention is a molecule produced by hybridization and/or linkage of at least two fragments or
15 elements to one another.

"Linkage"-- The "linkage" of two or more nucleic acid molecules to one another according to the present invention refers to any reaction that results in formation of a covalent bond between two nucleic acid molecules that were not covalently attached to one another prior to the linkage reaction. Preferably, the linkage is accomplished either
20 by splicing or by ligation. Alternatively, linkage may be accomplished indirectly, for example by replication of molecule pairs (or clusters) held together by ligation but including one or more nicks. Linkage may occur *in vitro* or *in vivo*.

"Overhang"- An "overhang", according to the present invention, is a single-stranded region of nucleic acid extending from a double-stranded region. Preferred overhangs are at least one nucleotide long. Particularly preferred overhangs are at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 20, or 25 nucleotides long. In some preferred
5 embodiments of the invention, the overhangs are comprised of at least one, preferably at least 2, 3, 4, 5, or more RNA residues; in other preferred embodiments the overhangs are comprised of DNA. In some embodiments of the invention, overhangs may comprise RNA elements that include functional intronic sequences.

"Portion"-- A "portion" of a nucleic acid molecule or polypeptide molecule, as
10 that term is used herein, is any piece that is shorter in length than the entire molecule. Preferably, a portion has a length sufficient to be characteristic of the full length molecule. For nucleic acid molecules, preferred portions are usually at least about 3-5 residues in length, more preferably at least about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, or
15 100 residues in length. For polypeptide molecules, preferred portions are typically at least about 2-5 residues in length, more preferably at least about 7, 10, 15, 20, 25, 30, or 40 residues in length.

"Primer"-- The term "primer", as used herein, refers to a polynucleotide molecule that is characterized by an ability to be extended against a template nucleic acid stand, so that a polynucleotide strand whose sequence is complementary to that of at least a portion
20 of the template strand, is produced linked to the primer. Preferred primers are at least approximately 5-10 nt long; particularly preferred primers are at least about 15 nt long. In many preferred embodiments, primers preferably have a length within the range of about 18-30 nt, preferably longer than approximately 20 nt.

Description of Certain Preferred Embodiments of the Invention

As described above, the present invention recognizes that vectors need not be provided as intact, discrete molecules, but rather can be provided as fragments that contain all or part of particular desired sequence elements. The invention provides techniques and reagents for the assembly of vectors (and/or inserts) through the linkage of such fragments. Certain preferred embodiments of this invention are described in more detail below.

Vector Elements

As will be appreciated by those of ordinary skill in the art, any desired nucleic acid sequence can be considered a vector element according to the present invention. Practitioners will be aware of their own needs and desires in terms of vector functions and attributes, and will readily be able to select appropriate sequences for use as vector elements. Nonetheless, certain types of sequence elements are already well established as useful in the field of vector construction. For example, Invitrogen Corporation, one of the larger distributors of molecular biology reagents, provides on its web site (www.invitrogen.com) a page entitled "Anatomy of a Vector" that lists the following categories of vector elements: promoters, inducible elements, transcriptional termination sequences, origins of DNA replication, affinity purification tags, multiple cloning sites/polylinkers, and selectable markers. The contents of this site, as they were presented on July 19, 2000, are included herein as Appendix B.

Replication elements

As described above, any sequence that operates to ensure replication of vector sequences in a selected host system constitutes a replication element. A variety of replication elements are already available in the art, and have been employed in commonly-available vector systems (see, for example, Ausubel et al., *Current Protocols in Molecular Biology*, Section II, Unit 1.5.1-1.5.17, John Wiley & Sons, 1998, the entire contents of which are incorporated herein by reference).

It will be appreciated by those of ordinary skill in the art that it is often desirable to construct a vector containing more than one replication element. For example, if it is desired that the same vector be able to replicate in more than one host cell type (e.g., in both bacterial cells and mammalian cells), then the vector should be designed to include replication elements that operate in each relevant cell type. On the other hand, it is also known that certain replication elements are incompatible with one another in a given cell type. It is generally desirable not to include incompatible elements in a single construct unless fragmentation of the construct in the host cell is desired.

Available replication elements that are known to operate in *E. coli*, the most commonly employed bacterium in molecular biology, include both high copy (so-called "relaxed control") elements such as pMB1 (100-300 copies/cell; Bolivar et al., *Gene* 2:95, 1977), ColE1 (>15 copies/cell; Kahn et al., *Method. Enzymol.* 68:268, 1979) and p15A (about 15 copies/cell; Chang et al., *J. Bacteriol.* 134:1141, 1978) and low copy (so-called "stringent control") elements such as pSC101 (about 6 copies/cell; Stoker et al., *Gene* 18:335, 1982), F (1 to 2 copies/cell; Kahn et al., *Method. Enzymol.* 68:268, 1979), and RK2 (2-4 copies/cell; Kahn et al., *Method. Enzymol.* 68:268, 1979). The R1 (low

copy at 30 °C and high copy above 35 °C; Uhlin et al., *Gene* 22:225, 1983) replicon also operates in *E. coli*, as do various phage origins of replication including λ *dv* (Jackson et al., *Proc. Natl. Acad. Sci. USA* 69:2904,1972), m13, f1, etc..

Replication elements that are known to operate in bacteria other than *E. coli* include RK2 and RSF1010, which have been shown, unlike ColE1, to have relatively broad host-ranges. In some cases, it may be desirable (or necessary) to introduce vectors into bacterial host cells through a mating process, in which case sequence elements encoding certain trans-acting factors (e.g., the *tra* or *mob* genes) may be required, as may be the cis-acting *oriT* site.

There are two primary categories of replication elements known to operate in yeast cells, centromeres and the 2 μ replicon. Of course, since DNA can readily be targeted for integration in yeast cells, it is not always necessary for a vector to be used in yeast cells to include an origin of replication that is active in those cells. Sequences that target integration of the vector into other replicating nucleic acid molecules are sufficient to constitute a replication element according to the present invention in those circumstances.

Several viral origins of replication, such as simian virus 40 [SV40], bovine papilloma virus [BPV], and Epstein Barr Virus [EBV], *oris* are known to operate in mammalian cells (sometimes requiring the presence of additional viral genes) and therefore can be employed as mammalian replication elements according to the present invention. Alternatively, sequences sufficient to target integration of a vector into another nucleic acid molecule (e.g., a chromosome or virus) capable of replicating in the mammalian cell can be employed. Targeted homologous recombination has been

demonstrated to work effectively in mammalian cells, so that regions of homologous gene sequence can operate as replication elements according to the invention.

Analogously, sequence elements of the Cre recombinase system can be employed to direct integration of vector sequences in mammalian systems (see, for example,

- 5 Fukushima et al., *Proc. Natl. Acad. Sci. USA*, 1992).

Viral origins of replication such as the baculovirus origin are known to operate in insect cells and can be employed as replication elements according to the present invention, as can other sequences, such as P-element sequences, that enable integration of vector sequences into other replication-competent nucleic acids.

10 In certain embodiments of the invention, it will be desirable to provide a particular replication element in two parts, on two different fragments, so that hybrid molecules will only replicate if they contain properly ligated fragments (see, for example, Figures 3, 4, and 6). In other embodiments, replication elements are provided intact on a single vector fragment (see, for example, Figures 2, 5, and 7-9).

15 *Vector detection elements*

A wide variety of sequences are available that allow host cells containing vector to be distinguished from host cells that do not contain vector. There are two basic categories of such elements: those that contain a selectable marker (i.e., one that imparts
20 a growth advantage to vector-containing cells under certain conditions) and those that contain a detectable marker. A wide variety of such markers is available, for use in different cell types.

09100001-072001
5 The most commonly employed selectable markers utilized in bacterial systems are those that confer resistance to antibiotics such as ampicillin, chloramphenicol, kanamycin, and tetracycline. Similarly, selectable markers commonly utilized in insect and/or mammalian cells include those that confer resistance to zeocin, neomycin, blasticidin, or hygromycin. The DHFR gene, which confers the ability to grow in the absence of exogenous purines (and also confers resistance to methotrexate, can also be used as a selectable marker in a range of cell types including mammalian cells. Also, cytosine deaminase can be used as a selectable marker under conditions that require cells to convert cytosine to uracil for growth. Other selectable markers useful in mammalian
10 cells include, for example, hygromycin- β -phosphotransferase (HPH), puromycin-N-acetyl transferase (PAC), thymidine kinase (TK), and xanthine-guanine phosphoribosyltransferase (XGPRT).

The most commonly employed selectable markers utilized in yeast cells include those that confer the ability to grow in the absence of a given nutrient such as uracil,
15 tryptophan, histidine, leucine, lysine, etc.

Preferred detectable markers for use in accordance with the present invention include genes encoding proteins that produce detectable products. Commonly employed detectable markers include, for example, the β -galactosidase gene, the green fluorescence protein gene, the horse radish peroxidase gene, the nitric oxide synthetase gene, the
20 chloramphenicol acetyl transferase gene, the luciferase gene, etc.

Those of ordinary skill in the art will readily appreciate that most or all of these vector detection elements can alternatively be employed as insert detection elements. For example, Figures 3-5 depict inventive reactions in which vector fragments are designed

so that, if they become linked to one another, a vector detection element is created. On the other hand, if an insert fragment becomes linked between them, the vector detection element is not created. Thus, constructs containing the insert fragment and those not containing the fragment can readily be distinguished from one another.

5 Similarly, those of ordinary skill in the art will appreciate that it will often be desirable to design vector and/or insert fragments so that a vector detection element is only created if the fragments become linked together in the desired arrangement. Figure 6, for example, depicts a particular embodiment of the invention in which this strategy was employed to simplify hybrid construct production according to the present invention.

10 It should be noted that one advantage of the present invention is that it renders the insert detection strategies described in the previous two paragraphs particularly practicable. The inventive modular approach to vector assembly, and particularly the inventive employment of cloning technologies that do not require restriction digestion, removes the need for a polylinker in order to introduce insert sequences into a vector.

15 Since polylinkers add unnatural sequences, their location in the middle of a detectable or selectable gene typically disrupted the gene activity, so that it was not possible to use reverse selection or detection to assay for insert insertion. By contrast, the inventive technologies allow the seamless union of insert and vector sequences, making feasible the use of these convenient screens and selections.

Expression elements

As will be appreciated by those of ordinary skill in the art, one of the most common uses of vector systems in molecular biology is to arrange for expression of insert

sequences in a host cell of interest. Any sequence that participates in directing or regulating expression of a linked sequence can be an expression element according to the present invention. A wide variety of such sequences are known in the art; certain examples are discussed in more detail below.

5

PROMOTER: Promoters are the regions of DNA that are responsible for establishing the initiation site for transcription. A variety of different promoters, operative in different systems, have been defined and characterized. Different promoters may direct expression of linked sequences at different levels. Furthermore, some
10 promoters are constitutively active, while others can have their activity modulated through adjustment of the experimental conditions. Some promoters are active in only particular cell types, where as others are ubiquitously expressed.

Preferred promoters known to be active in bacterial cells include, for example, P_{BAD}, P_L, P_R, *lack*, *tack*, *trc*, *spa lacUV5*, T3, T7, T7 LAC, SP6, etc.; preferred promoters
15 known to be active in yeast cells include, for example p*GAL1*, p*AOX1*, p*ADH*, etc.; preferred promoters known to be active in insect cells include, for example, the MT, Ac5, and polyhedrin promoters, etc; preferred promoters known to be active in mammalian cells include, for example, P_{ΔHSP}, P_{SG}, P_{CMV}, P_{EF-1α}, P_{SV40}, P_{RSV}, P_{PGK}, P_{MMTV}, P_{MC1} etc.

ENHANCERS/TRANSCRIPTIONAL REGULATORS: Regulator sequences that operate to
20 stimulate or repress transcription from a given promoter in certain cell types or under certain conditions can often be combined with any of a variety of different promoters to create a transcription control element with useful characteristic. The universe of known regulatory sequences operative in different organisms is very large. Particularly preferred

elements that are commonly used in vector systems include, for example, the lac operon, the λ cI site, the tet operon, lexA sites, Gal4 sites, the SV40 enhancer, the MMTV enhancer, etc. Those of ordinary skill in the art will immediately recognize the huge range of alternative sequences that could be employed in the practice of the present invention. Experiments to define additional such sequences, operative in the context of any particular experiment, are routine.

TRANSCRIPTION TERMINATOR: Although not required, it is sometimes desirable to include in an expression vector sequences that will terminate transcription of relevant sequences at a selected point. Without such termination signals, it may be possible for RNA polymerases, at least under some circumstances, to transcribe indefinitely around a circular construct. A variety of different transcriptional termination sequences have been identified; the one most commonly used in vector applications is probably the SV40 terminator. Alternatively or additionally, 3'-end formation signals, such as polyadenylation sites, may be employed.

SPLICING SIGNALS: In certain circumstances, it may be desirable to include in inventive expression vectors signals that can direct splicing of transcripts encoded by insert sequences. For example, if a vector includes a promoter and exonic sequences including a splice donor site, then insert sequences containing a splice acceptor site can be expressed and translated. In certain embodiments of the invention, it might be desirable to provide a collection of vectors or vector fragments (3) that contain the splice acceptor site in all three possible frames, so as to ensure in-frame fusions of insert sequences in one version of the vector, regardless of whether information about the insert sequence is available.

TRANSLATION START: Often, if expression of an insert that does not include 5' sequences (or is not known to include such sequences) is desired, it will be useful to include translation start sequences. The consensus translation start sequence, known as the Kozak sequence, will provide the strongest translation initiation signal, but in most cases a single ATG reasonably positioned with respect to the start of the transcript will suffice.

TRANSLATION STOP: Expression vectors designed to express insert sequences that may be lacking their natural 3' ends often benefit from the inclusion of translation stop sequences. As with the translation start and splicing sequences, families of vectors can be prepared containing the relevant sequences in all three possible frames so that knowledge of the insert sequence is not required. Alternatively, a single vector could be employed but families of insert fragments can be prepared with additional (or fewer) nucleotides on one or both ends.

15 *Gene fusions*

As those of ordinary skill in the art will be aware, a variety of vector systems have been engineered to generate gene fusions between insert sequences and a reporter gene in the vector backbone. Such fusions are useful, for example, to detect expression patterns of the insert sequences, or to detect expression control elements that may be present in the insert sequences. Gene fusions may also allow a researcher to track the expression products of the fused gene.

Particularly preferred detectable genes for use in gene fusion applications include, for example, LacZ, chloramphenicol acetyl transferase (CAT), green fluorescence protein (GFP), luciferase, horse radish peroxidase (HRP), etc.

5 *Fusion proteins*

One version of gene fusions that is particularly commonly employed in vector systems is fusions that generate fusion proteins with a desirable characteristic. As will be appreciated, it will often be desirable to provide families of vectors or vector fragments that allow C-terminal, N-terminal, or internal fusions, and also that allow fusions in all possible frames, preferably without knowledge of insert sequence.

For example, a variety of sequence elements are available that encode polypeptides that, when fused to a polypeptide encoded by an insert sequence, allow that polypeptide to be readily purified. Particularly preferred purification tags include, for example, (His)₆, thioredoxin, glutathione-S-transferase, streptavidin, staphylococcal protein A (which interacts strongly with IgG; Amersham Pharmacia Biotech, Piscataway, NJ), etc.

Also available are a variety of sequence elements encoding detectable moieties, such as epitopes for which high-specificity antibodies are available, that can be useful in the detection of an expression fusion protein. Examples of such detectable epitopes include, for example, Xpress™, *c-myc*, CA25, thioredoxin, V5, HA, calmodulin binding peptide (CBP), Aag, etc.

In some cases, it is desirable to remove the protein tags created by fusion of encoding insert sequences with encoding vector sequences. Sequence elements encoding

polypeptide cleavage elements (e.g., by furin, enterokinase, thrombin, factor X1, PreScission, etc.) are particularly useful in such applications.

Other useful sequence elements for the production of fusion proteins are ones that encode targeting moieties, such as secretion signals (e.g., BiP for insect cells, human placental alkaline phosphatase or human growth hormone for mammalian cells, protein A for bacterial cells, etc.) or other elements, that direct the fusion product to a particular cellular location. Examples of such targeting sequences include, for instance, yeast AgA2 sequences that target the fusion protein to the cell surface, VP22 fusions that target to the mammalian nucleus, pRLT3-NLS, COXVIII signal, etc.

Polylinkers

One virtually ubiquitous element in most commercially-available vectors today is a so-called "polylinker" or "multiple cloning site". In certain embodiments of the invention, it may be desirable to include vector fragments containing such elements in linkage reactions. However, in many embodiments, it will be desirable to create fragments and/or hybrid molecules without employing the use of restriction enzymes. As techniques for such restriction-free nucleic acid manipulation become more accepted, the need for polylinkers in inventive vectors and reactions will diminish.

Other elements

Those of ordinary skill in the art will readily appreciate that any of a variety of other sequence elements may be included in vector fragments according to the present invention. The foregoing has been intended to provide merely a sampling of certain

examples of sequence elements that are currently commonly found in vector sequences.

One of the advantages of the present invention is that, by providing techniques and reagents that allow the ready production of specifically designed vectors through the assembly of prepared fragments, it is expected that the invention will also help

5 researchers expand the range of sequence elements utilized in vector applications.

Insert Elements

As will be apparent to those of ordinary skill in the art, any nucleic acid sequence may be employed as an insert element according to the present invention. A researcher
10 may choose any sequence or sequences s/he likes to be linked to vector sequences. Also, more than one insert element may be employed. Furthermore, each insert element may be provided as a single insert fragment, or may be distributed over multiple insert fragments that will be linked together in series in the final hybrid product. In certain
15 embodiments of the invention, part or all of a given insert element may even be prepared as a single fragment that also includes part or all of one or more vector elements. Any collection of contiguous insert sequences is considered a single insert element for the purpose of the present invention.

Those of ordinary skill in the art will recognize that the classification of particular sequences as "insert elements" as compared with "vector elements" is not critical to the
20 invention. In fact, the inventive recognition that a "vector" need not be a single discrete molecular entity in a sense renders such distinctions arbitrary. Nonetheless, both the concept and the terminology of a "vector backbone" and an "insert" are well established

in molecular biology and therefore can be useful for the purposes of clarity and communication.

Preparation and Linkage of Fragments

5 In general, any method may be used to prepare fragments for hybridization and/or linkage according to the present invention. However, it is preferred that, for each hybrid molecule to be assembled, at least one fragment is prepared without the use of restriction enzymes, and preferably without the use of any endonuclease.

10 In certain preferred embodiments of the invention, fragments are prepared in a form that allows them to be linked together by ligation. In other embodiments, fragments are prepared in a manner that allows them to be linked together by splicing. In particular, U.S. patent applications USSN 08/814,412, USSN 09/399,593, USSN 09/225,990, and PCT/US00/0189, and US Patents 5,498,531 and 5,780,272, each of which is incorporated herein by reference, contain thorough descriptions of methods and strategies useful in the
15 preparation of nucleic acid (RNA or DNA) fragments that contain flanking intronic sequences and can be linked to one another by trans- or cis-splicing. In yet other embodiments, fragments are prepared in a form that allows topoisomerase-mediated linkage.

20 Often, it will be desirable to prepare fragments so that, for each linkage reaction to be performed in the assembly of a hybrid molecule, the fragments are designed to associate with one another in only one way and to produce only a single major linkage product. For example, fragments may be prepared so that each has single-stranded overhangs on one or both ends, and only fragments that are to be adjacent to one another

in a hybrid molecule have complementary overhangs. Alternatively or additionally, fragments may be engineered to include intronic elements that are only compatible with the intronic elements on adjacent fragments. Such "directed linkage" (i.e., linkage in only one arrangement) of fragments discussed above is particularly desirable where multiple fragments (i.e., three or more, preferably four or more, and more preferably five or more) are to be linked together in a single linkage reaction. For linkage reactions containing small numbers of fragments (2 or 3), directed linkage can be assured by controlling the phosphorylation state of the relevant fragment ends.

In other preferred embodiments of the invention, it may be desirable to prepare fragments so that they can become linked to one another in any of a variety of different ways. This phenomenon is referred to herein as "linkage degeneracy". In such embodiments, a single linkage reaction can generate a "library" of different hybrid molecules that can subsequently be distinguished and/or separated from one another as desired.

In yet other preferred embodiments of the invention, fragments can be designed for directed ligation as described above, but then multiple alternative versions of each particular fragment can be provided in the same linkage reaction so that, once again, a library of hybrid molecules is produced in a single linkage reaction. This phenomenon is referred to herein as "selection degeneracy". For example, fragments A, B, and C can be designed and prepared so that they can only be linked to one another in the arrangement ABC (which can be a linear or a circular arrangement). If multiple different A fragments (e.g., A1, A2, A3, . . . An), multiple different B fragments, and/or multiple different C fragments are employed in a single linkage reaction, then a library of different hybrid

molecules, each having an ABC structure, will be produced in that reaction (e.g., A1B17C3, A1B1C1, A1B2C1, etc.). Those of ordinary skill in the art will readily appreciate that the different versions of the A fragment need not bear any relationship to one another other than being designed to be link only to a B fragment, etc. Alternatively, 5 each version of a given fragment could, for example, contain different varieties of the same vector element(s) or element portion(s) (e.g., different drug resistance genes).

Still other preferred embodiments combine the two kinds of degeneracy discussed above, so that a single linkage reaction may create a library of hybrid molecules in which both the arrangement and selection of fragments is varied.

10 According to the present invention, particularly preferred fragments for use in accordance with the present invention contain one or more single-stranded overhangs available for hybridization with complementary overhangs on other fragments. It is most preferred that such overhang-containing fragments be prepared without the use of restriction enzymes. It is particularly preferred that such fragments be prepared using 15 RNA-Overhang Cloning (ROC) or DNA-Overhang Cloning (DOC), as described for example in USSN 09/225,990; PCT US00/00189; and USSN 09/478,263, each of which is incorporated herein by reference in its entirety (see also Examples 5-8).

Once a hybrid molecule is created by hybridization or linkage of vector and/or insert fragments, it may be replicated by any available *in vitro* or *in vivo* mechanism. In 20 certain preferred embodiments of the invention, hybridization or linkage reactions themselves, or isolated or purified hybrids prepared from such reactions (e.g., by gel electrophoresis), may be directly transformed or transfected into host cells (or otherwise introduced into a host system). In some cases, it may be desirable to perform one or

more manipulations prior to introducing a hybrid molecule into a host cell. For example, linkage of two fragments created using some embodiments the ROC methodology will produce a hybrid molecule that includes some regions of double-stranded RNA that may not be stable inside certain host cells. Accordingly, it may be desirable to perform at least a single round of DNA replication of such a hybrid prior to introducing it into a cell. Other circumstances in which such additional manipulations (e.g., nick repair, etc.) are desirable will be apparent to one of ordinary skill in the art.

Kits

As discussed herein, one aspect of the present invention comprises the recognition that vectors are comprised of modular elements and can be assembled from individually prepared fragments. One part of this recognition includes the realization that vector fragments can be provided as isolated cassettes, ready to be assembled by a user or a manufacturer.

In one embodiment of the invention, a variety of different possible vector elements is offered to a user who selects particular pieces of interest. Fragments that together comprise these pieces are then prepared and are provided to the user for assembly into a vector. Optionally, reagents for performing the assembly (e.g., ligase if the fragments are prepared with overhangs amenable to linkage by ligation; splicing reagents if the fragments are prepared for linkage by splicing; etc.). Alternatively, the fragments may be linked together into a "designer vector" before being provided to the user.

09910354-072001

In other embodiments of the invention, kits are provided that contain multiple optional fragments, each of which contains a selected vector element or elements, or fragment(s) thereof, so that a user can readily assemble any of a variety of different vectors by mixing different collections of fragments together in linkage reactions. For example, a bacterial expression vector kit could be provided that contains (a) a first collection of first fragments, each of which contains the pTac promoter and also contains a portion of an antibiotic resistance gene, where different fragments in the collection contain portions of different antibiotic resistance genes; (b) a second collection of second fragments, each of which contains the remainder of one of the antibiotic resistance genes and also contains the ColE1 origin of replication; and (c) ligation reagents. A user could then select particular first and second fragments that, when ligated with his or her chosen insert fragment(s), would create a hybrid containing a chosen antibiotic resistance gene and the insert element under control of the pTac promoter. Those of ordinary skill in the art will immediately recognize the infinite variety of related other kits that could alternatively or additionally be provided.

The inventive recognition of vector modularity also provides a new perspective on valuable reagents, and systems for providing such reagents to users. For example, in addition to kits as discussed above, reagent providers could prepare catalogs or menus (either paper or electronic) from which users can select particular desired vector elements or fragments. In certain preferred embodiments of the invention, the catalog or menu is presented on a World Wide Web site that the user can access and through which the user can place an order. In other embodiments a paper form is provided, or information about telephone contact is provided. As discussed above, selected fragments may be provided

to the user as isolated fragments, as fragment collections, as linked pieces (e.g., complete vectors), as kits (e.g., including linkage reagents, purification reagents, amplification reagents, instructions for use and/or other relevant materials), or in any other desirable form. The invention therefore provides, in addition to the various techniques and reagents discussed herein, new methods of doing business in the area of molecular biology reagents.

Hybrid Molecules

As discussed herein, the techniques and reagents provided by the present invention allow the ready assembly of any of a variety of hybrid molecules, generated by hybridization and/or linkage of vector and/or insert fragments. In some embodiments of the invention, a vector is assembled from vector fragments (via one or more than one linkage reactions) prior to linkage of vector sequences with insert sequences. In other embodiments, assembly of the complete, final hybrid product is accomplished in a single linkage reaction. In other embodiments, one or more linkage fragments is/are linked to one or more vector fragments in a first linkage reaction, and one or more additional linkage reactions are subsequently performed to add additional vector and/or insert fragments. Each and every hybrid molecule produced in such a linkage reaction is encompassed within the scope of the present invention.

Examples

Example 1

Assembly of a λ Vector/Insert Hybrid

Figure 1 presents an inventive reaction for the assembly of a hybrid molecule containing two λ phage arms (a λ cloning vector) separated by a chosen insert. As is well known, λ vectors are particularly useful for the cloning of relatively large (up to about 50 kB) fragments. The insert-containing hybrids can be packaged (typically through the use of helper phages) into phage heads *in vitro*. Although the efficiency of packaging can be relatively low (around 10%), the subsequent efficiency of genome transfer into bacteria through infection is close to 100% (see, for example, Ausubel et al., *Current Protocols in Molecular Biology*, Unit 1.10, Current Protocols, 1987, the entire contents of which are incorporated herein by reference).

Example 2

Assembly of a Bacterial Vector/Insert Hybrid

Figure 2 presents an inventive reaction for the assembly of a hybrid molecule containing a bacterial origin of replication, an antibiotic resistance gene, and a chosen insert. The hybrid molecule is assembled by linkage of three fragments, each of which contains a single element. Preferably, the fragments are prepared to have complementary overhangs selected to provide for directional ligation. Alternatively, the indicated element in each fragment may be flanked by intronic components that direct appropriate trans-splicing reactions *in vivo* or *in vitro*.

Example 3

Assembly of a Bacterial Vector/Insert Hybrid in Which the Insert Disrupts a Detectable Element

Element

29 of 41

The inventive reaction depicted in Figure 3 differs from that shown in Figure 2 (and discussed above in Example 2) in at least two ways. First, the two vector fragments employed in the reaction of Figure 3 each contain a part of the bacterial origin of replication, so that only hybrid molecules in which these two fragments are properly
5 linked together will be able to replicate in bacteria. Also, each vector fragment contains a portion of the ampicillin resistance gene. If a hybrid is assembled that does not include an insert, the ampicillin resistance gene will be re-created (unless some mutation occurs) and bacteria containing the resulting hybrid will be resistant to both tetracycline and ampicillin. By contrast, the ampicillin gene will not be re-created in hybrid that do
10 contain the insert. Thus, bacteria containing complete hybrids will be distinguishable from those containing partial hybrids that lack insert because those containing complete hybrids will be resistant to tetracycline but not ampicillin, whereas those containing partial hybrids will be resistant to both.

The strategy depicted in Figure 3 is particularly useful for fragments that do not
15 have directionally specific ends. For example, if blunt-ended fragments are to be employed, or if the both ends of the insert fragment (and both ampicillin fragment ends) contain identical overhangs, the ability to identify desirable hybrids from the universe of possible hybrids is particularly useful.

The strategy depicted in Figure 4 is analogous to that depicted in Figure 3 except
20 that hybrids containing insert are distinguishable from those lacking insert on the basis of a blue/white screen rather than a growth/no growth screen.

The strategy depicted in Figure 5 is also similar, except that linkage of the vector fragments is not required to create a functional origin of replication. For this strategy, it

is generally preferred that at least the vector fragments be engineered for directional linkage , so that they can only be linked to one another in a single orientation.

Example 4

5 Assembly of a Hybrid Bacterial Expression Vector/Insert Construct by 4-Way Ligation

The inventive strategy depicted in Figure 6 shows simultaneous linkage of three different vector fragments with an insert fragment. A hybrid vector molecule containing both an origin of replication and an ampicillin resistance gene can only be assembled through proper linkage of the three vector fragments. Thus, selection strategies can be employed to identify desirable hybrid molecules. Such molecules can then be screened for expression of the insert in order to identify those that are complete as compared with those that contain only vector sequences.

Example 5

15 Assembly of a Hybrid Bacterial Vector/Insert Molecule Using DOC

The inventive scheme depicted in Figure 9 was carried out as follows. Vector fragments were amplified from the pET 19b vector (Novagen, Madison, WI.) using the following primers (lower case letters indicate RNA residues; upper case letters indicate DNA residues): EV-1 (5'- cauGGTATATCTCCTTCTTAAAG; SEQ ID NO:1), EV-2 (5'- cucATGACCAAAATCCCTTAAC; SEQ ID NO:2), EV-3 (5'- gagATTATCAAAAAGGATCTTC; SEQ ID NO:3), and EV-4 (5'- uaaCTAGCATAACCCCTTGG; SEQ ID NO:4). EV-1 and EV-2 were used together to generate vector fragment 1, containing the bacterial origin of replication, the LacI gene,

and the pT7 promoter; EV-3 and EV-4 were used together to generate vector fragment 2, containing the Amp gene.

In a separate DOC reaction, an insert fragment containing the *Lac Z* gene was amplified from the pBluescript II SK (-) vector (Stratagene, La Jolla, CA), with primers
5' Lac Z (5'-augACCATGATTACGCCAACG; SEQ ID NO:5) and 3' Lac Z (5'-
uuaCAATTTCATTCGCCATTC; SEQ ID NO:6). 100 µl PCR reactions contained 5 ng
of template DNA, 1x cloned PFU buffer (Stratagene, La Jolla, CA), 1mM MgSO₄, 200
µM of each dNTP, 1.45 U cloned PFU (Stratagene), 1.25 U PFU Turbo™ polymerase and
50 pM of each primer. Reactions were performed in a Robocycler (Stratagene, La Jolla,
CA) as follows: 1 cycle 95°C, 5 min; 53°C, 3 min; 72°C, 6 min (10 min for vector
fragment 1); 30 cycles, 95°C, 1 min; 53°C, 1 min; 72°C, 3 min (8 min for vector
fragment 1) ; and 1 cycle 72°C 10 min.

Products of the PCR reactions were separated on a 1% agarose gel, and purified
using the GENECLAN II kit (Vista, CA.). 12 µl of each purified fragment was placed
separately in 1x first strand buffer (Life Technologies, Rockville, MD.) with 10 mM
DTT, 5 mM of each dNTP, and 200 U M-MLV (Life Technologies). Reactions were
incubated for 20 min at 42°C. Reactions were then placed at 70°C for 10 min to heat kill
the enzyme.

Primer ribonucleotides were removed from the PCR products by hydrolysis with
NaOH. 6 µL of 1 N NaOH were added to each reaction, and the mixtures were incubated
for 30 min at 45°C. 6 µl of 1 N HCL, 4 µl of 10x ligase buffer (USB, Cleveland, OH),
and 10 U of T4 PNK (USB) were then added. Reactions were incubated at 37°C for 30

min. Phosphorylated fragments were combined in equimolar amounts (approximately 50 ng) and ligated with 10 U of T4 DNA ligase (USB) at 25°C for 2 hrs. 5 µl of the ligation reaction was then transformed into *E.coli*.

5

Example 6

Assembly of Hybrid Vector/Insert Molecules Using ROC with Internal Terminators

We prepared hybrid vectors containing an origin of replication (Ori) fragment and a kanamycin resistance gene (KAN) fragment, by amplifying each fragment with primers that contained one or more residues not copied by the DNA polymerase utilized in the reaction (i.e., "terminator" residues). The Ori fragment was amplified from pET19b (Novagen, Madison, WI); the KAN fragment was amplified from pCR 2.1 (Invitrogen, Carlsbad, CA). Figure 11 shows the various primers used and fragments generated. As will be seen, some reactions generated a 2400 bp ori fragment; others generated an 824 bp fragment, (denoted "Ori(s)" because it is smaller). The smaller fragment, Ori(s), lacks an 11 pb direct repeat that can create a deletion hotspot when it is present.

PCR reaction cycling, product annealing and *E. Coli* transformation were performed as described in Examples 7 and 8.

20

Example 7

Assembly of a Hybrid Vector/Insert Molecule Using ROC with Single Nucleotide Terminators

The vector/insert hybrid molecule depicted in Figure 10 was generated as follows. The ori-containing vector fragment was amplified from pET 19b (Novagen, Madison, WI) using primers (lower case letters indicate RNA residues; upper case letters indicate DNA residues) 5'OST (5'- CTGCTAAGTGAGcucGACAGATCGCTGAGATAGGTGC; SEQ

5 ID NO:5) and 1N3' Ori(s)(5'-

AAGCTTGCTAAGTAaggCGTTTTTCCATAGGCTCCG; SEQ ID NO:6)

The vector fragment containing the Kanamycin resistance gene was amplified from pCR2.1 Topo (Invitrogen, Carlsbad, CA) using primers 1NT5'KAN (5' CTACCTAGCAAGCTuCTATCTGGACAAGGGAACG; SEQ ID NO:7) and T7 3'KAN (5'CCCTATAGTGAGTCGTATTaAGCGAAAACCTCAAGGATC; SEQ ID NO:8).

The insert fragment containing the luciferase gene was amplified from pGI II basic (Promega, WI) using primers tCS1 (5' TTAATACGACTCACTATAGGGATGGAAGACGCCAAAAACATA; SEQ ID NO: 9) and tCS8 (5'- GAGTCACTTAGCAgTTACAATTTGGACTTCCGCC; SEQ ID NO: 10).

Each 100 µl reaction contained 50 pM of each primer, 1x cloned *Pfu* Buffer (10 mM (NH₄)₂SO₄, 20 mM Tris (pH 8.8), 2 mM Mg SO₄, 10 mM KCE, 0.1% Triton x-100 and 0.1 mg/me Bovine serum Albumin), 1mM additional mg SO₄, 0.3 mM each dNTP, 5- 10 ng template DNA and 1.25-1.85 units of both cloned *Pfu* and *Pfu* Turbo polymerases (strategies, La Jolla, CA). The Ori fragment was amplified in a reaction involving (1) one cycle of 95° for 3 min; 46-60° for 2 min; (2) 35 cycles of 95° for 30 sec; 48-60° for 30 sec; and 72° for 3 min; and (3) one cycle of 95° for 30 sec; 48-60° for 30 sec; and 72° for

8 min. The KAN and LUC fragments were amplified in similar reactions except that the 35 cycles contained a 4.5 min 72° step.

PCR products generated in these reactions were gel purified using the Qiaquick gel extraction kit (Qiagen, Valencia, Ca.). Approximately 80 ng of each fragment was
5 combined in a 20 µl reaction volume. Two (2) µl 10x USB ligation Buffer (660mM Tris-HCL (pH7.6), 66 mM MgCl₄, 100 mM DTT, 660 µM ATP) (USB, Cleveland, OH) was then added, to make a 1x reaction mix. The reaction was heated to 65° C for 8 minutes, and then slow cooled for 20 minutes (to 35-40° C) to allow the fragments to anneal. Samples were spun and allowed incubate another 15 minutes at room temperature.

The annealing reaction was precipitated by adding 100 µl of 100% ethanol,
10 followed by a 15 minute incubation at -80° C, and a 70% and 100% wash. Electrocompetent DH5α cells were transformed using a Biorad E. coli pulser (Biorad, Hercules, CA). Five (5) µl of each annealing reaction was combined with 40 µl of
15 Elexctromax DH5α-E cells (Life technologies, Rockville, MD) Individual clones generated in this experiment were isolated, restriction mapped, and sequenced; all junctions were correct.

Those of ordinary skill in the art will appreciate that, as with Example 6, the ROC
technique described in this Example utilizes primers containing internal ribonucleotide
residues (in one case, 3 residues were used; in other cases only one) flanked by DNA
20 residues. The overhangs created in these ROC PCR reactions, therefore, have only a
single "ribo" residue; other overhang residues are DNA. In separate experiments, we
have demonstrated that any individual ribonucleotide (i.e., rA, rG, rU, or rC) can act
effectively to block extension of a complimentary strand by an appropriate DNA

polymerase, so that overhangs are created (see, for example, Example 6). We have also
owed that single 3'-Omethyl residues are similarly effective. Primers containing 3'-
Omethyl residues can often be synthesized more easily (e.g., due to higher coupling
efficiencies) than those containing inbanucleotides, and will generally be more stable, so
5 that they are preferred for many applications.

Example 8

Streamlined Cloning

Inventive modular vector fragments may be prepared, annealed together, and
10 transformed into host cells, without enzymatic ligation. For example, we assembled a
two-fragment vector, by preparing one fragment (KAN) containing the kanamycin
resistance gene, and one fragment (Ori) containing an origin of replication.

Specifically, two 100 μ l PCR reactions were performed to amplify each of the
two components of the vector. Each reaction contained 50 pM of each primer, 1x Cloned
15 *Pfu* Buffer (10 mM $(\text{NH}_4)_2\text{SO}_4$, 20 mM Tris (pH 8.8), 2 mM MgSO_4 , 10 mM KCl, 0.1%
Triton X-100 and 0.1 mg/ml bovine serum albumin), 1 mM additional MgSO_4 , 0.3 mM
of each dNTP, 5-10 ng of plasmid template and 1.25-1.85 units of both cloned *Pfu* and
Pfu Turbo polymerases (Stratagene, La Jolla, Ca.).

The following chimeric RNA/DNA primers were purchased from Oligo's Etc. (
20 Willsonville, OR): (ribonucleotides are in lower case)

INT 5'KAN-CTACCTAGCAAGCTuCTATCTGGACAAGGAAAAACG (SEQ ID
NO:11)

INT 3'KAN-GAGCTCACTTAGCA^aGGCGAAAACTCTCAAGGA (SEQ ID NO:12)

INT5' Ori- TTGCTAAGTGAGCUcGACAGATCGCTGAGATAGGTGC (SEQ ID NO:13)

IN3' Ori(s)-AAGCTTGCTAAGTA_gGGCGTTTTTCCATAGGCTCCG (SEQ ID

- NO:14). Primers INT 5'KAN and INT 3'KAN were used to amplify the Kan fragment from pCR 2.1 Topo (Invitrogen, Carlsbad, CA). Primers INT5' Ori and IN3' Ori(s) were used to amplify the Ori fragment from pET 19b (Novagen, Madison, WI). The following cycles were performed: one cycle of 95°, 3', 48-60°, 2', 72°, 8'; followed by 35 cycles of 95°, 30sec, 48-60°, 30 sec, 72°, 3' for Ori fragment, 4.5' for Kan and Luc fragments. A final cycle with an 8' 72° step was performed in all cases.

Approximately 80 ng of each fragment (5 µl each) produced in the PCR reactions was combined and mixed. 5 µl of this reaction was then transformed into 100 µl of chemically competent DH5α cells. Positive clones were isolated and mapped; sequence junctions appear to be correct.

Other Embodiments

Those of ordinary skill in the art will appreciate that the foregoing represents certain preferred embodiments of the invention, but is not intended to limit the spirit or scope of the following claims.